

Please replace the paragraph beginning at page 32, line 21 with the following rewritten paragraph:

a7
10025947
2-16-5200T
--The PC3 cell line was maintained in RPMI supplemented with 10% fetal calf serum and antibiotics. Cells were suspended in 0.12% soft agar in complete medium and plated (2,000 cells per well) in different drug concentrations onto a 0.4% agarose underlayer in 24-well plates. Plating cells on agarose underlayers supports the proliferation only of the transformed cells, ensuring that the growth signal stems from the malignant component of the tumor.--


Please replace the paragraph beginning at page 37, line 24 with the following rewritten paragraph:

a8
10025947
2-16-5200T
--Cells were treated with test compounds and CFTR immunoprecipitated as described in Bradbury et al., *Am. J. Physiol.* 276, L659 - 668 (1999). Briefly, treated cells were lysed in buffer containing 1% TRITON X-100 and various protease inhibitors. Soluble material was immunoprecipitated using both R domain and C-terminal monoclonal antibodies. Immunoprecipitated CFTR was then subject to *in vitro* phosphorylation using camp-dependent PKA catalytic subunit and [γ -32P]ATP, followed by resolution on SDS-PAGE gels. After fixation, the gels were dried and processed for autoradiography and phosphor image analysis. Quantitation of B and C bands was performed on a BioRad personal fix image analysis station.--

In the claims:

Please amend claims 1, 20, 21, 22 and 43 as follows:

--1. A compound of formula (I):



A is a cyclic moiety selected from the group consisting of C₃₋₁₄ cycloalkyl, 3-14 membered heterocycloalkyl, C₄₋₁₄ cycloalkenyl, 3-14 membered heterocycloalkenyl, aryl, and heteroaryl; the cyclic moiety being optionally substituted with alkyl, alkenyl, alkynyl, alkoxy, hydroxyl, hydroxylalkyl, halo, haloalkyl, amino, alkylcarbonyloxy, alkyloxycarbonyl, alkylcarbonyl, alkylsulfonylamino, aminosulfonyl, or alkylsulfonyl;

each of X¹ and X², independently, is O or S;

each of Y¹ and Y², independently, is -CH₂-, -O-, -S-, -N(R^a)-, -N(R^a)-C(O)-O-, -O-C(O)-N(R^a)-, -N(R^a)-C(O)-N(R^b)-, -O-C(O)-O-, or a bond; each of R^a and R^b, independently, being hydrogen, alkyl, alkenyl, alkynyl, alkoxy, hydroxylalkyl, hydroxyl, or haloalkyl;

L is a straight C₃₋₁₂ hydrocarbon chain optionally containing at least one double bond, at least one triple bond, or at least one double bond and one triple bond; said hydrocarbon chain being optionally substituted with C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, hydroxyl, halo, amino, nitro, cyano, C₃₋₅ cycloalkyl, 3-5 membered heterocycloalkyl, monocyclic aryl, 5-6 membered heteroaryl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ alkyloxycarbonyl, C₁₋₄ alkylcarbonyl, or formyl; and further being optionally interrupted by -O-, -N(R^c)-, -N(R^c)-C(O)-O-, -O-C(O)-N(R^c)-, -N(R^c)-C(O)-N(R^d)-, or -O-C(O)-O-; each of R^c and R^d, independently, being hydrogen, alkyl, alkenyl, alkynyl, alkoxy, hydroxylalkyl, hydroxyl, or haloalkyl; provided that when L contains two or more double bonds, the double bonds are not adjacent to each other; that when L contains three double bonds, said hydrocarbon chain is substituted with C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, hydroxyl, halo, amino, nitro, cyano, C₃₋₅ cycloalkyl, 3-5 membered heterocycloalkyl, monocyclic aryl, 5-6 membered heteroaryl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ alkyloxycarbonyl, C₁₋₄ alkylcarbonyl, or formyl; and further provided that when L contains less than 6 carbon atoms in the hydrocarbon chain and A is C₁₋₄ alkyl phenyl or unsubstituted phenyl, Y¹ is not a bond;


or a salt thereof.--

--20. The compound of claim 1, said compound being 4-chloro-5-phenyl-2,4-pentadienoic acid, 5-(4-dimethylaminophenyl)-2,4-pentadienoic acid, 5-(2-furyl)-2,4-

pentadienoic acid, 5-phenyl-2-en-4-yn-pentanoic acid, or 8-phenyl-3,5,7-octatrienoic acid.--

--21. The compound of claim 1, said compound being 8-phenyl-3,5,7-octatrienoic acid.--

--22. A compound of formula (I):



wherein

A is a cyclic moiety selected from the group consisting of aryl and heteroaryl; the cyclic moiety being optionally substituted with alkyl, alkenyl, alkynyl, alkoxy, hydroxylalkyl, or amino;

each of X¹ and X², independently, is O or S;

each of Y¹ and Y², independently, is -CH₂-, -O-, -S-, -N(R^a)-, -N(R^a)-C(O)-O-, -O-C(O)-N(R^a)-, -N(R^a)-C(O)-N(R^b)-, -O-C(O)-O-, or a bond; each of R^a and R^b, independently, being hydrogen, alkyl, hydroxylalkyl, or haloalkyl;

L is a straight C₃₋₁₂ hydrocarbon chain optionally containing at least one double bond, at least one triple bond, or at least one double bond and one triple bond; said hydrocarbon chain being optionally substituted with C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, or amino, and further optionally interrupted by -O- or -N(R^c)-, where R^c is hydrogen, alkyl, hydroxylalkyl, or haloalkyl; provided that when L contains two or more double bonds, the double bonds are not adjacent to each other; that when L contains three double bonds, said hydrocarbon chain is substituted with C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, hydroxyl, halo, amino, nitro, cyano, C₃₋₅ cycloalkyl, 3-5 membered heterocycloalkyl, monocyclic aryl, 5-6 membered heteroaryl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ alkyloxycarbonyl, C₁₋₄ alkylcarbonyl, or formyl; and further provided that when L contains less than 6 carbon atoms in the hydrocarbon chain and A is C₁₋₄ alkyl phenyl or unsubstituted phenyl, Y¹ is not a bond;
or a salt thereof.--

Applicant : Hsuan-Yin Hargest et al.
Serial No. : 09/812,940
Filed : March 27, 2001
Page : 7

Attorney's Docket No.: 12938-003002

a¹¹
--43. The compound of claim 40, wherein L is an unsaturated C₄₋₈ hydrocarbon chain containing at least one double bond in trans configuration, said unsaturated hydrocarbon chain being optionally substituted with C₁₋₂ alkyl, C₁₋₂ alkoxy, hydroxyl, -NH₂, -NH(C₁₋₂ alkyl), or -N(C₁₋₂ alkyl)₂.--

10025947-122601
T0923T 4652001